

Charged particle multiplicity in Pb-Pb collisions from NA50 experiment

F. Prino for the NA50 collaboration

*International Workshop on
Particle Multiplicity in Relativistic Heavy Ion Collisions
Bari, 17-19 June 2004*

Particle production in nuclear collisions

- **Multiplicity = number of particles produced in the collision**
 - ➔ 80-90% of the produced charged particles are pions
 - ➔ Related to the centrality of the collision
 - ➔ Related to the entropy of the system created in the collision
- **Multicollision models:**
 - ➔ Nucleus-nucleus collis.= superposition of nucleon-nucleon collis.

Hard processes

Large momentum transfer

Small distance

- ✓ Interactions at partonic level
- ✓ Scale like the number of elementary collisions (N_{coll})

Soft processes

Small momentum transfer


Large distance

- ✓ Interactions at the baryon level
- ✓ Scale like the number of participant nucleons (N_{part})

Particle momenta distributions

Particle momenta decomposed

Transverse momentum (p_T)
 Longitudinal momentum (p_L)



- Rapidity variable

$$y = \frac{1}{2} \cdot \ln \left(\frac{E + p_L}{E - p_L} \right)$$

- Pseudorapidity variable


$$\eta = -\ln \left[\tan \left(\frac{\vartheta}{2} \right) \right] = \frac{1}{2} \cdot \ln \left(\frac{|\mathbf{p}| + p_L}{|\mathbf{p}| - p_L} \right)$$

- $\eta \approx y$ for large momenta

➔ dN/dy ($dN/d\eta$) distributions carry information about energy density, longitudinal expansion and "stopping power"

NA50: experimental setup

Study of muon pair production in Pb-Pb collisions



Trigger

➔ DIMU

2 muon tracks

➔ MB

Non zero energy deposit in the ZDC

● Pb beam

➔ 1998: 158 GeV/nucleon

➔ 1999: 40 GeV/nucleon

● Beam detectors

● Active target

➔ Up to 7 Pb subtargets + Cherenkov counters

● Centrality detectors

➔ EM Calorimeter ($1.1 < \eta_{\text{lab}} < 2.3$)


➔ **Multiplicity Detector** ($1.1 < \eta_{\text{lab}} < 4.2$)

➔ Zero Degree Calorimeter ($\eta_{\text{lab}} > 6.3$)

● Muon spectrometer ($2.7 < \eta_{\text{lab}} < 3.9$)

➔ Magnet+MWPC+hodoscopes

The Multiplicity Detector (MD)



Silicon microstrip detector measuring the number and the angular distribution of charged particles produced in the collision

- 2 Planes (MD1, MD2)


- each plane made of 2 layers (up/down)

- 36 azimuthal sectors ($\Delta\phi=10^\circ$)

- 192 radial strips ($\Delta\eta=0.02$)

- 6912 strips in each plane

- Only 128 innermost strips used in this analysis



dN/d η distributions vs. centrality (I)

- Data from special low-intensity runs


<i>Pb beam energy (GeV/nucleon)</i>	<i>Target</i>	<i>Distance target-MD1 (cm)</i>	<i>Target thickness</i>	<i># of events analyzed</i>
158	Pb	11.65	3 mm	48000
158	Pb	9.15	1 mm	18000
40	Pb	12.55	3 mm	35000

- Analysis method

➤ Data selection:

- ✓ Interaction trigger
- ✓ Pile-up rejection
- ✓ Upstream interaction rejection
- ✓ Diagonal cut on the E_T - E_{ZDC} correlation
- ✓ MD based target identification

- Statistical method based on matching pairs of hits on MD1 and MD2





dN/dη distributions vs. centrality (II)

→ Centrality interval definition at 158 GeV/c:



- ✓ 2 independent centrality variables (E_T and E_{ZDC})
- ✓ Intervals expressed in terms of fraction of total inelastic cross section

$$N_{\text{int}} = N_{Pb} \cdot P_{\text{int}} = N_{Pb} \cdot \left(1 - e^{-L_T/\lambda_{INT}}\right) \quad \text{with} \quad \lambda_{INT} = \frac{A_{TARG}}{\rho_{Pb} N_A \sigma_{inel}} = 3.98_{-0.27}^{+0.21} \text{ cm}$$




$dN/d\eta$ distributions vs. centrality (III)

- Calculation of raw $dN_{\text{ch}}/d\eta$
 - ✓ Cluster (group of contiguous strips firing together) correction
 - Cluster size distribution not reproduced by a VENUS+GEANT simulation
 - Dedicated MC, aimed at reproducing cluster size distribution observed in data
 - ✓ Performed separately in each η bin ($\Delta\eta=0.15$) and in each centrality class



$dN/d\eta$ distributions vs. centrality (III)



- Calculation of primary $dN_{ch}/d\eta$.
 - ✓ Subtraction of the delta electron contribution (from GEANT).
 - Max. 5% of the occupancy in the most peripheral bin.
 - ✓ Correction with secondary/primary ratio from VENUS+GEANT simulation.
 - VENUS+GEANT data reconstructed with same method as experimental data.
 - 1.2 – 1.8 correction factor.
 - *Do not depend on centrality.*
 - *Depend on target thickness, target position, particular MD plane.*
 - Unstable particles (K_0 , Λ and hyperons) decays are already considered in VENUS, and therefore their decay products are defined as primary particles.
- Systematic error estimation
 - ✓ **8% systematic error** on primary charged multiplicity



$dN/d\eta$ distributions vs. centrality (III)

Agreement between MD1 and MD2

- ➔ Average between detector planes
- ➔ Wide η coverage




Excellent agreement between primary $dN_{ch}/d\eta$ with 2 different target thicknesses and positions


- ➔ Average between different thicknesses
- ➔ Wider η coverage

$dN_{ch}/d\eta$ distributions at 158 GeV

E_T centrality selection



E_{ZDC} centrality selection




Distributions fitted with Gaussians to extract


- Midrapidity value η
- Gaussian width
- $dN_{ch}/d\eta$ at the peak

Midrapidity value at 158 GeV /c

E_T centrality selection



E_{ZDC} centrality selection




Midrapidity visible in the $dN/d\eta$ distributions:


- No reflection around midrapidity needed
- η_{peak} extracted from fit compatible with VENUS prediction ($\eta_{\text{peak}}=3.1$)

Gaussian width at 158 GeV /c

E_T centrality selection



E_{ZDC} centrality selection




Gaussian width decreases with centrality:


- stopping power effect
- decreasing contribution of protons from target and projectile fragmentation

$dN_{ch}/d\eta$ $_{\text{max}}$ at 158 GeV /c

E_T centrality selection



E_{ZDC} centrality selection



$dN/d\eta$ at the peak scales linearly with E_T and E_{ZDC}

→ no saturation or enhancement observed



$dN_{ch}/d\eta$ distributions at 40 GeV /c (I)

ZDC worse performance
at such a low energy


Only E_T based **centrality selection**

Larger (10%) systematic error

- ZDC based quality cuts not performed
 - larger E_T tail beyond the knee
 - E_T resolution not well defined





$dN_{ch}/d\eta$ distributions at 40 GeV /c (II)




Peak position
(VENUS prediction):
 $\eta_{max} \approx 2.47$

Gaussian width smaller
than at 158 GeV,
decreasing with increasing
centrality



Number of charged
particles in the
more central bin
 ≈ 2 times smaller
than at 158 GeV

Gaussian width vs. energy



- Available phase space in rapidity increases with \sqrt{s}
- Fit with the simple scaling law: $\sigma_\eta = a + b \cdot \ln \sqrt{s}$
- Same \sqrt{s} dependence for all data

Evaluation of N_{part} and N_{coll} (I)

• Glauber model calculations

➔ Physical inputs:

✓ Woods-Saxon density for Pb nucleus (2pF) →




✓ $\sigma_{\text{in}} = 30 \text{ mb}$

➔ Numerical calculation of:

✓ Interaction probability, N_{part} , N_{coll} ... vs. impact parameter b

$$\rho = \frac{\rho_0}{1 + e^{(r-r_0)/C}}$$

$\rho_0 = 0.16 \text{ fm}^{-3}$ (green arrow)
 $r_0 = 6.624 \text{ fm}$ (red arrow)
 $C = 0.549 \text{ fm}$ (blue arrow)



Evaluation of N_{part} and N_{coll} (II)

• E_T and E_{ZDC} parametrization

✓ $E_T \propto$ number of participants

✓ $E_{ZDC} \propto$ number of projectile spectators



• q, w, α and δ form fit to MB spectra

$$E_T = q \cdot N_{part}$$

$$\sigma^2_{E_T} = w \cdot q^2 \cdot N_{part}$$


$$E_{ZDC} = E_{beam} \left(A_{Pb} - \frac{N_{part}}{2} \right) + \alpha N_{part}$$

$$\sigma^2_{E_{ZDC}} = \left(\beta \sqrt{E_{ZDC}} + \gamma E_{ZDC} \right)^2 + \delta^2$$



Evaluation of N_{part} and N_{coll} (III)

- Calculation of $\langle N_{part} \rangle$ and $\langle N_{coll} \rangle$ in each centrality class
 - ➔ From distributions of N_{part} and N_{coll} in the E_T and E_{ZDC} intervals
 - ➔ Smearing effects due to calorimeter resolution included



Class	% of c.s.	$E_{ZDC}^{min}-E_{ZDC}^{max}$ (GeV)	$\langle N_{part} \rangle$	RMS N_{part}	$\langle N_{coll} \rangle$	RMS N_{coll}
1	0-5	0-9385	354	22	802	66
2	5-10	9385-13150	294	23	634	63
3	10-15	13150-16490	246	25	501	64
4	15-20	16490-19180	205	26	395	65
5	20-25	19180-21475	173	28	316	66
6	25-35	21475-24790	129	35	214	74

Class	% of c.s.	$E_T^{min}-E_T^{max}$ (GeV)	$\langle N_{part} \rangle$	RMS N_{part}	$\langle N_{coll} \rangle$	RMS N_{coll}
1	0-5	87.2-140.	352	25	796	73
2	5-10	71.5-87.2	294	26	632	72
3	10-15	58.7-71.5	245	23	498	61
4	15-20	48.9-58.7	203	20	392	52
5	20-25	40.9-48.9	169	19	309	44
6	25-35	29.6-40.9	127	20	213	45



Charged particle scaling at 158 GeV

- Fit with the power law: $\left. \frac{dN}{d\eta} \right|_{\max} \propto N_{part}^{\alpha}$
 - $\alpha = 1.00 \pm 0.01 \pm 0.04$
 - ✓ $\alpha = 1.05-1.08$ using a VENUS calculation of N_{part}
 - ✓ $\alpha = 1.02$ with $N_{part} = 2 \cdot 208 \cdot (1 - E_{ZDC}/E_{BEAM})$


- Fit with the power law: $\left. \frac{dN}{d\eta} \right|_{\max} \propto N_{coll}^{\beta}$
 - $\beta = 0.75 \pm 0.02$

- Fit with the law: $\left. \frac{dN}{d\eta} \right|_{\max} \propto A \cdot N_{part} + B \cdot N_{coll}$
 - B compatible with zero

- Conclusions:
 - N_{part} describes the centrality dependence of particle production
 - Hard processes play a negligible role at this energy



Charged particle scaling at 40 GeV




Fit with the power law:

➤ $\alpha = 1.02 \pm 0.02 \pm 0.06$

Conclusions:

- As expected, no important hard process contribution at this energy
- Same N_{part} dependence at 158 and 40 GeV
- Soft processes account well for particle production at SPS energies

Yield per participant pair vs. centrality



Yield per participant pair:

$$\frac{dN_{ch}/d\eta|_{\max}}{0.5 N_{part}}$$


→ *Only statistical error on $dN/d\eta$ + error on N_{part} shown in plot*

→ *Flat behaviour reflects the linear dependence of $dN/d\eta|_{\max}$ on N_{part}*

Beam energy (A·GeV/c)	\sqrt{s} (GeV)	$\frac{dN_{ch}/d\eta _{\max}}{0.5 N_{part}}$
40	8.77	$1.18 \pm 0.03 \pm 0.15$
158	17.3	$2.49 \pm 0.03 \pm 0.20$

Comparison with other experiments

- Conversion from $dN/d\eta|_{lab}$ to dN/dy and subsequently to $dN/d\eta|_{cm}$ done assuming:
 - At 158 GeV/c ($\sqrt{s}=17.3$ GeV): pions, protons and kaons relative yields from NA49
 - At 40 GeV/c ($\sqrt{s}=8.77$ GeV): pions, protons and kaons relative yields from VENUS 4.12




$$\frac{dN_{ch}}{d\mathbf{p}_T d\eta} = \sqrt{1 - \frac{m^2}{m_T^2 \cosh^2 y}} \frac{dN_{ch}}{d\mathbf{p}_T dy}$$

most central bin

\sqrt{s} (GeV)	$\frac{dN_{ch}/d\eta _{\max}}{0.5 N_{part}} (LAB)$	$\frac{dN_{ch}/d\eta _{\max}}{0.5 N_{part}} (CMS)$
8.77	$1.18 \pm 0.03 \pm 0.15$	$0.97 \pm 0.03 \pm 0.14$
17.3	$2.49 \pm 0.03 \pm 0.20$	$2.14 \pm 0.03 \pm 0.17$


Yield per participant pair vs. \sqrt{s}

**158 GeV/c result:
50% higher than fit to pp inelastic
20% higher than fit to pp NSD**



**40 GeV/c result:
in agreement with
fit to pp inelastic**

Integrated yield per participant pair



Heavy ion data does not follow the e^+e^- trend over the whole energy range:

- below pp and e^+e^- data at AGS energy
- cross through pp data at SPS energy
- joins e^+e^- data above top SPS energy

Conclusions

- Particle pseudorapidity distribution vs. centrality measured by NA50 experiment.
 - At 158 GeV/c with 2 independent centrality estimators (E_T , E_{ZDC}).
 - At 40 GeV/c with 1 centrality estimator (E_T).
- Use of 4 detector planes + 2 different target positions.
 - Cross check of analysis procedure.
 - Wide η coverage (\rightarrow no reflection around midrapidity needed).
- Gaussian width:
 - Decreases with centrality (stopping power effect).
 - Increases logarithmically with \sqrt{s} (phase space effect).
- Glauber calculation of N_{part} and N_{coll} :
 - Linear dependence of $dN/d\eta|_{\text{max}}$ on N_{part} .
 - No important role of hard interactions (N_{coll}) at both energies.
- Yield per participant pair.
 - ✓ At 40 GeV/c ($\sqrt{s}=8.77$ GeV) compatible with fit to nucleon-nucleon interactions.
 - ✓ At 158 GeV/c ($\sqrt{s}=17.3$ GeV) not compatible with fit to nucleon-nucleon interactions.
 - Steep increase of particle yield in central Pb-Pb collisions between 40 and 158 GeV/c not described by the simple energy scaling observed in nucleon nucleon collisions.