

NA38/NA50 experiments



# Recent results on intermediate mass dimuon production

L.Capelli

Institut de Physique Nucléaire de Lyon, France



# Outline

- **introduction**
- **data analysis**
  - ✓ p-A
  - ✓ S-U and Pb-Pb
- **comparison with models**
  - ✓ charm enhancement
  - ✓ D-mesons rescattering
  - ✓ thermal dimuons
- **conclusions**




# Introduction

- ❖ NA50 has shown that the p-A dimuon mass spectra in the mass range 1.5 to 2.5 GeV/c<sup>2</sup> is correctly reproduced by a superposition of DY and D $\bar{D}$  dimuons
- ❖ a linear extrapolation of p-A sources to nucleus-nucleus collisions underestimate data [[Euro.Phys.J.C14\(2000\)443](#)]
- ❖ new development on this subject using a 4-dimensional unfolding method [[NIM.A405\(1998\)139](#)]
- ❖ NA50 apparatus
  - ➔ detect opposite sign ( $\mu^+\mu^-$ ) muon pairs
  - ➔ centrality detection
- ❖ record the like sign pairs  $\mu^+\mu^+$  and  $\mu^-\mu^-$ 
  - ➔ combinatorial background



# NA50 Experiment





# Unfolding method

❖ detector effects → **acceptance** and **resolution**

acceptance  $A(x)$      $x$  : set of kinematical variables describing  
resolution  $S(x'|x)$     the dimuon →  $M, p_T, Y_{cm}, \cos(\theta_{cs})$

$$D(x') = \int S(x'|x) A(x) \Phi(x) dx$$


⇒ extract the physical distribution  $\Phi$  from the measured one  $D$

❖ 4-D unfolding method [**NIM A405(1998)139**]

- ✓ based on image restoration methods extended to 4-D
- ✓ accounts for detector correlation
- ✓ preserve physics correlations
- ✓ no need to assume specific shapes for distribution
- ✓ iterative method



# Dimuon mass spectrum



❖ Known sources :

- ✓ Drell-Yan :  $q\bar{q} \rightarrow \mu^+\mu^-$
- ✓ resonances decay :  $J/\psi, \psi' \rightarrow \mu^+\mu^-$
- ✓ charmed meson (and baryon) decays :  $D \rightarrow \mu^+\mu^-$



# Mass spectra analysis

fit data in the mass range  $1.6 < M < 8.0$  GeV/c<sup>2</sup> assuming

$$\frac{dN}{dM} = n_1 \frac{dN^{D\bar{D}}}{dM} + n_2 \frac{dN^{DY}}{dM} + n_3 \frac{dN^{\psi}}{dM} + n_4 \frac{dN^{\psi'}}{dM}$$

- ❖ gaussian shapes for the  $J/\psi$  and  $\psi'$  resonances
- ❖ shapes of  $DY$  and  $D\bar{D}$  obtained from **PYTHIA** 6.1 with :
  - ✓ c quark mass  $\Rightarrow m_c = 1.5$  GeV/c<sup>2</sup>
  - ✓ intrinsic transverse momentum
$$\sigma_{k_T}^{DY} = 0.8 \text{ GeV/c [NA51 pp collisions]}$$
$$\sigma_{k_T}^{D\bar{D}} = 1.0 \text{ GeV/c [Eur.Phys.J.C1(98)123]}$$
  - ✓ **MRS A** set of PDF's


- ❖ 7 parameters fit




# p-A results - 450 GeV

measured data unfolded in the following kinematical domain :

$$M > 1.6 \text{ GeV}/c^2, -0.2 < Y_{\text{cm}} < 0.4 \text{ and } -0.3 < \cos(\theta_{\text{cs}}) < 0.3$$




$$\left. \frac{D\bar{D}}{DY} \right|_{\text{pp},450} = 4.13 \pm 0.15$$


in  $1.6 < M < 2.5 \text{ GeV}/c^2$



# Open charm cross section

➤ the open charm cross section @ 450 GeV is deduced in the following way :

$$\sigma_{c\bar{c}}^{450} = \sigma_{IMR}^{DY} \times \left. \frac{D\bar{D}}{DY} \right|_{pp,450} \times (\text{phase space factor}) \times \frac{1}{BR(D \rightarrow \mu)^2}$$



➤ the value is compatible with other direct measurements




# Ion mass spectra


- ❖ analysis as a function of centrality based on electromagnetic transverse energy  $E_T$  :
  - ✓ 5 bins for S-U
  - ✓ 7 bins for Pb-Pb
- ❖ data unfolded in the following kinematical domain
$$\begin{cases} M > 1.6 \text{ GeV}/c^2 \\ 0.2 < Y_{\text{cm}} < 0.8 \\ -0.3 < \cos(\theta_{\text{cs}}) < 0.3 \end{cases}$$
- ❖ for ion collisions, the DY and  $D\bar{D}$  processes are extrapolated linearly from NN yields, as expected for hard processes
- ❖ NN open charm and DY cross sections have been deduced from the p-A 450 GeV/c value using the  $\sqrt{s}$ -dependence given by PYTHIA
- ❖ the isospin correction has been taken into account for DY



# Pb-Pb mass spectra




➔ in the IMR, data are **higher** than the expected sources



# data/expected sources

❖ quantify the difference between data and expected sources as a function of centrality

➤ plot data/expected sources vs  $N_{\text{part}}$  in  $1.6 < M < 2.5 \text{ GeV}/c^2$



❖ the IMR **excess increases** as a function of **centrality**




# Charm enhancement

- ❖ hypothesis : excess behaves as open charm  
[C.Y. Wong and Z.Q. Wang, Phys.Lett.**B367**(96)50]
- fit the IMR ion mass spectra with a superposition of DY and  $D\bar{D}$  and extract the ratio  $(D\bar{D}/DY)_{measured}$
- calculate the expected ratio  $(D\bar{D}/DY)_{expected}$  from p-A
- ❖ plot the enhancement factor **E** as a function of centrality


$$E = \frac{\left(D\bar{D}/DY\right)_{measured}}{\left(D\bar{D}/DY\right)_{expected}}$$



# Charm enhancement




- ✓ charm-like enhancement : factor  $\sim 3$  in **central Pb-Pb** with respect to p-A
- ✓ **linear increase with  $N_{part}$**




# D-mesons rescattering

- ❖ several theoretical models have been proposed to explain the observed IMR charm excess
- ❖ Z. Lin and X.N. Wang [[Phys.Lett.B444\(98\)245](#)] associate the observed excess to D-mesons rescattering in nuclear matter which leads to an enhancement in the limited phase space of the NA50 experiment
- ❖ D and  $\bar{D}$  rescattering is described by a thermal distribution depending on a temperature parameter T



# D-mesons rescattering

- ❖ the enhancement factor in NA50 phase space is calculated as the ratio of the number of dimuons observed at temperature  $T$  and at  $T=0$




with MRS A  
and  
 $m_c = 1.5 \text{ GeV}/c^2$




# D-mesons rescattering

- ❖ from the experimental value of the enhancement the corresponding temperature can be obtained for each of the centrality bins


$$E_{\text{data}}(N_{\text{part}}) \text{ and } E_{\text{model}}(T)$$




- ❖ the shape of the dimuon mass distribution from  $D\bar{D}$  decays is then calculated with the corresponding temperature



# D-mesons rescattering



❖ the mass shape calculated with this model fails to reproduce the IMR mass spectra in central Pb-Pb collisions




# Thermal dimuons

❖ model developed by Rapp and Shuryak  
[Phys.Lett.**B473**(2000)13]

- $\mu\mu$  yield based on  $q\bar{q}$  annihilation rate
- integration over space-time history
- central collisions only
- parameters :
  - fireball lifetime : 14 fm/c
  - initial temperature :  $T_i = 192$  MeV
- explicit introduction of a QGP phase
  - critical temperature :  $T_c = 175$  MeV



# Thermal dimuons



→ the IMR excess can be well accounted for by **thermal radiation** when combined with **DY** and **open charm**



# Conclusions

- ❖ the  $\sigma_{c\bar{c}}$  cross section extracted from the p-A data agrees with direct measurements of other experiments
- ❖ the ion data are in excess of the DY+D $\bar{D}$  superposition extrapolated from p-A
- ❖ this excess increases linearly with  $N_{\text{part}}$
- ❖ the mass distribution cannot be reproduced by a model assuming D and  $\bar{D}$  rescattering
- ❖ two possible explanations of the observed excess :
  - ✓ the data can be described under the hypothesis of an enhancement of charm production
  - ✓ the central Pb-Pb mass distribution can be reasonably well reproduced by the thermal model